## Predicting the Pedestal in a Burning Plasma: Progress and Research Needs

#### **Philip Snyder**

Theory and Computational Science Department General Atomics San Diego CA, USA

Presented at US Burning Plasma Organization Web Seminar

21 February 2019







J.W. Hughes, O. Meneghini, T.H. Osborne, W. Solomon, H.R. Wilson, N. Aiba, D. Battaglia, E.A. Belli, M. Beurskens, K. Burrell, J. Candy, J. Canik, C.S. Chang,
A. Diallo, D. Dickinson, M. Dorf, B. Dudson, M. Dunne, D. Eldon, M. Fenstermacher, J. Ferron, A. Garofalo, B. Grierson, R.J. Groebner, W. Guttenfelder, D. Hatch, A. Hubbard, G.T.A. Huijsmans, Y. Kamada, A. Kirk, M. Komm, C. Konz, M. Kotschenreuther,
B. LaBombard, F. Laggner, L. Lao, A.W. Leonard, M. Leyland, C. Maggi, R. Maingi, W. Meyer, R. Nazikian, D. Orlov, N. Oyama, J.M. Park, S. Parker, C. Paz-Soldan, S. Saarelma,
G.M. Staebler, A. Turnbull, H. Urano, E. Viezzer, J.R. Walk, E. Wang, E. Wolfrum, X. Xu, and the ITPA Pedestal Group





#### Fundamental Challenge: Fusion Conditions in Core Compatible with Edge/Materials



- Core plasma 10x hotter than the core of the sun
- Need increase of ~100x in temperature, ~1000x in pressure to reach fusion conditions in core





#### Fundamental Challenge: Fusion Conditions in Core Compatible with Edge/Materials



- Physics of turbulent transport and large scale MHD instabilities constrain average pressure and temperature gradients
  - Large normalized size ( $aI_pB_t \sim a^2B_pB_t$ ) needed, potentially expensive

a=minor radius,  $I_p$ =plasma current= $B_p$ \*circumference,  $B_t$ =toroidal magnetic field



#### Fundamental Challenge: Fusion Conditions in Core Compatible with Edge/Materials



- Physics of turbulent transport and large scale MHD instabilities constrain average pressure and temperature gradients
  - Large normalized size (al\_pB\_t ~ a^2B\_pB\_t) needed, potentially expensive
    - a=minor radius, I<sub>p</sub>=plasma current=B<sub>p</sub>\*circumference, B<sub>t</sub>=toroidal magnetic field
  - Ideal solution: Suppress turbulence and build sharp gradients across outermost part of confined plasma
    - Broad profiles for high global pressure limit, large fusion volume



### Sometimes Nature Comes Through: H-Mode and Pedestal

- In 1982 ASDEX reported a new regime named H-mode
  - Later found on numerous tokamaks
- Factor of ~2 improvement in confinement and stored energy
  - Bursty events called Edge Localized Modes (ELMs)
- Characterized by suppression of turbulence and high gradients in outer few % of confined plasma



# The H-mode Transport Barrier is Enabled by Shear in the Edge Rotation Profile

- Driven by auxiliary heating, a spontaneous transition to a higher confinement state
- Shear in perpendicular rotation (Er × B) suppresses turbulent transport
  - Shears long wavelength turbulence of Lmode, reducing transport, gradients rise
  - Labeled "E<sub>r</sub> well"
- Detailed physics of "L-H" transition complex (key research need) – here focus on physics controlling structure of the pedestal



### H-mode Produces a Steep Edge Pressure Gradient

- Narrow edge layer of steep pressure gradient
- High pressure core plasma rests on this "Edge Pedestal"





### The Pedestal is Narrow but Impactful

- Narrow edge layer of steep pressure gradient
- High pressure core plasma rests on this "Edge Pedestal"
- Can have >10x increase in T, and >40x increase in p across this layer
  - Typically larger relative increase than core





### Fusion Performance Rests Upon the Pedestal

- Future burning plasmas rely on maintaining high pedestal pressure P<sub>fusion</sub> p<sup>2</sup><sub>ped</sub>
- For ITG limited cores, T<sub>i,ped</sub> plays a key role in global confinement
  - On existing devices, optimizing fusion performance a combination of a T<sub>i,ped</sub> and p<sub>ped</sub> optimization
  - However, considering all microinstabilities in core, pressure broadening for global MHD stability, and i-e coupling at reactor scale, p<sub>ped</sub> generally most important

### **ITER Fusion Power**



[J. Kinsey, Nucl. Fusion 51 083001 (2011)]



# Outline: Pedestal Physics Key to Predicting and Optimizing the Tokamak

- The Pedestal: What it is and why it matters
  - Simultaneous improvement of confinement and stability
  - Predictive capability enables fusion power optimization ( $P_{fus} \sim p_{ped}^2$ )
- Rich physics and computational challenges
  - Overlap of scales, challenge to methods (L $\sim\lambda\sim\rho$ )

#### Physics approaches and experimental tests

- Gyrokinetics and neoclassical theory
- MHD and peeling-ballooning modes

#### • Predicting and optimizing the pedestal: The EPED model

- Development and testing
- Coupled core-pedestal prediction fusion optimization
- Super H-Mode and high fusion performance
- ITER predictions and future prospects



#### Outline: Pedestal Physics Key to Predicting and Optimizing the Tokamak

- The Pedestal: What it is and why it matters
  - Simultaneous improvement of confinement and stability
  - Predictive capability enables fusion power optimization ( $P_{fus} \sim p_{ped}^2$ )
- Rich physics and computational challenges
  - Overlap of scales, challenge to methods (L~ $\lambda$ ~ $\rho$ )
- Physics approaches and experimental tests
  - Gyrokinetics and neoclassical theory
  - MHD and peeling-ballooning modes
  - Diagnostics and dedicated experiments
- Predicting and optimizing the pedestal: The EPED model
  - Development and testing
  - Coupled core-pedestal prediction fusion optimization
- Super H-Mode and high fusion performance
- ITER predictions and future prospects



### Very Wide Range of Overlapping Scales in the Edge Barrier Region



Both time and spatial scales overlap, from microscopic all the way to global

• This wide range (6-7 orders of magnitude) is covered by a **single** equilibrium, key parameters vary by orders of magnitude across the pedestal



### Pedestal Physics Challenges Existing Paradigms



- Our field traditionally divided into stability (λ~ L >>p), transport (λ~p<< L) and source physics (λ=fluctuation scale, L=equilibrium scale, p=drift-/gyro- orbit scale)</li>
  - This separation can break down in the edge barrier
- Simulations focused on 3D collisional or 5D collisionless equations
  - Edge barrier is in general both highly collisional and highly collisionless
  - Perturbations can be large, potential problem for  $\delta f$
  - Electromagnetic perturbations (and 3D fields) and full geometry important



## Traditional Transport Theory Requires a Separation of Scales

- Fluctuation scale=λ
- Equilibrium scale=L (eg pressure gradient scale L<sub>p</sub>)
- Microscopic scale=p (toroidal or poloidal gyroradius)
- Standard transport theory allows ( $\lambda \sim \rho$ ), expands in  $\rho/L$

Leading order: gyrokinetic and neoclassical fluxes

Next order: evolution of equilibrium (L>> $\lambda \sim \rho$ )

Equilibrium scale macrostability (MHD) (L $\sim\lambda$ >> $\rho$ )

In the pedestal, fluctuation scale overlaps equilibrium and micro scales (L $\sim\lambda\sim\rho$ ), transport theory formally breaks down

- Can proceed using existing tools to develop physics insight, but must be cautious of limits (in particular the L>> $\lambda$  approximation can lead to arbitrarily large errors for ion scale modes)

### <u>Open issue</u>: Work on extended formulations or 6D simulations by many authors, but practical, rigorous formalism for pedestal remains a challenge

GENERAL ATOMICS

# Outline: Pedestal Physics Key to Predicting and Optimizing the Tokamak

- The Pedestal: What it is and why it matters
  - Simultaneous improvement of confinement and stability
  - Predictive capability enables fusion power optimization ( $P_{fus} \sim p_{ped}^2$ )
- Rich physics and computational challenges
  - Overlap of scales, challenge to methods (L~ $\lambda$ ~ $\rho$ )

#### Physics approaches and experimental tests

- Gyrokinetics and neoclassical theory
- MHD and peeling-ballooning modes
- Predicting and optimizing the pedestal: The EPED model
  - Development and testing
  - Coupled core-pedestal prediction fusion optimization
- Super H-Mode and high fusion performance
- ITER predictions and future prospects

16

#### The Fokker-Planck Equation Provides the Fundamental Theory for Plasma Equilibrium, Fluctuations, and Transport

### • $\left[\frac{\partial}{\partial t} + \vec{v} \cdot \nabla + \frac{z_a e}{m_a} \left(\vec{E} + \hat{\vec{E}}\right) \cdot \frac{\partial}{\partial \vec{v}} + \frac{z_a e}{m_a c} \vec{v} \times \left(\vec{B} + \hat{\vec{B}}\right) \cdot \frac{\partial}{\partial \vec{v}}\right] (f_a + \hat{f}_a) =$ $\sum_b C_{ab} (f_a + \hat{f}_a, f_b + \hat{f}_b) + S_a$ $f_a, \vec{E}, \vec{B} \rightarrow \text{ensemble-averaged}$ $\hat{f}_a, \hat{\vec{E}}, \hat{\vec{B}} \rightarrow \text{fluctuating}$

 Separate the FP equation into ensemble-averaged (A) and fluctuation (F) components.

• 
$$A = \left[\frac{\partial}{\partial t} + \vec{v} \cdot \nabla + \frac{z_a e}{m_a} \left(\vec{E} + \frac{\vec{v}}{c} \times \vec{B}\right) \cdot \frac{\partial}{\partial \vec{v}}\right] f_a - \langle C_a \rangle_{ens} - D_a - S_a = 0 \rightarrow \mathsf{DKE}$$

- $\mathbf{F} = \left[\frac{\partial}{\partial t} + \vec{v} \cdot \nabla + \frac{z_a e}{m_a} \left(\vec{E} + \frac{\vec{v}}{c} \times \vec{B}\right) \cdot \frac{\partial}{\partial \vec{v}}\right] \hat{f}_a + \frac{z_a e}{m_a} \left(\vec{E} + \frac{\vec{v}}{c} \times \vec{B}\right) \cdot \frac{\partial}{\partial \vec{v}} (f_a + \hat{f}_a) C_a + \langle C_a \rangle_{ens} + D_a = 0 \rightarrow \mathbf{GKE}$
- Use the drift-ordering to separate neoclassical and turbulent transport:  $\rho_* = \rho_i/L << 1$

Leading order, neoclassical and turbulence separate ( $D \Rightarrow 0$ ).



## Gyroaverages of the O( $\rho_*$ ) Ensemble-averaged and Fluctuating Equations Give the Drift-kinetic and Gyrokinetic Equations

• Drift-Kinetic Equation:

$$\oint \frac{d\xi}{2\pi} A_1 = 0:$$

$$f_{1a} = \tilde{f}_{1a} + \bar{f}_{1a}$$

gyroangle- gyroangledependent independent

The first-order (gyroangle-independent) ensemble-averaged distribution is determined by the DKE

$$\mathbf{v'}_{\parallel}\hat{b}\cdot\nabla\left(\bar{f}_{1a}-f_{0a}\frac{z_{a}e}{T_{a}}\Phi_{1}\right)-\sum_{b}C_{ab}^{L}\left(\bar{f}_{1a},\bar{f}_{1b}\right)=S_{neo}$$

• Gyrokinetic Equation:  $\oint \frac{d\xi}{2\pi} F_1 = 0: \ \hat{f}_{1a}(\vec{x}) = -\frac{z_a e}{T_a} \widehat{\Phi}_1 + h_a(\vec{x} - \vec{\rho})$ 

$$\frac{\partial h_a(R)}{\partial t} + \left(\vec{V}_0 + \mathbf{v'}_{\parallel}\hat{b} + \mathbf{v}_{da} - \frac{c}{B}\nabla\widehat{\Psi}_a \times \hat{b}\right) \cdot \nabla h_a - \sum_b C_{ab}^{GL}(\widehat{f}_{1a}, \widehat{f}_{1b}) = S_{turb}$$

The first-order fluctuating distribution in terms of the distribution of gyrocenters  $h_a(R)$  is determined by the GKE

18

### Neoclassical Bootstrap Current in Pedestal Validated Against Observations, Key for Instability Physics

 Pressure gradient gives rise to toroidal bootstrap current

$$j_b \propto rac{dp/dr}{B_{ heta} \left(1 + 0.9 \sqrt{v_e^*}
ight)}$$

- Efficient calculation with codes like NEO [Belli9,12] and NEOART [Peeters00], reduced models [eg Sauter99] and neural nets for very fast evaluation
- Experimental validation of pedestal bootstrap current,  $\pm$  20%
- Large bootstrap current reduces magnetic shear and can both stabilize and drive pedestal instabilities

Open issue: Higher order (finite orbit width) effects explored by many authors but theory complexity and exp't validation unresolved



## Neoclassical Ion Heat Transport Very Important in the Pedestal

- High resolution measurements and efficient calculations (eg NEO/NEOART) have confirmed important role for neoclasssical ion heat transport
  - In some cases can account for ~all inter-ELM ion heat flux [ASDEX-U, Viezzer16]
  - Even in cases with strong fluctuations, remains significant
- While neoclassical particle flux is small in a pure plasma, inward pinch of impurities very important, particularly with high-Z materials



#### Gyrofluid and Gyrokinetic Edge Simulations Identified Role of Electromagnetic Modes, KBM



- Electrostatic limit requires (at least) that: (a)  $\beta$  is small, (b) frequency small compared to shear Alfven frequency, (c) p' far from ideal ballooning limit (a<<1 or  $d\beta_p / d\psi_N <<1$ )
  - (c) is nearly always violated in the pedestal due to sharp gradients, and (b) can be violated as well (small k<sub>par</sub>, drift-Alfven modes) [Scott98, Snyder99]
- Kinetic Ballooning mode (KBM) goes unstable just below ideal ballooning mode threshold due to ion drift resonance
- Recent studies have assessed effects of multiple species, full collisions [Belli17]

GENERAL ATO

### Consideration of Non-local Effects Key for General Assessment of KBM



Open issue: Kinetic ballooning mode calculations with full non-locality (including kink term)

- Purely local (infinite n ballooning, flux tube GK) calculations predict "second stability" for KBM at low magnetic shear (high bootstrap current)
- Finite-n MHD studies find non-local effects close 2<sup>nd</sup> stability
  - kink term found to be important even at very high n
- Non-local gyrokinetic studies find similar behavior [Wan/Parker12, Saarelma/Dickinson17, Lin16]
  - Mode continues to have finite pressure gradient threshold even at low magnetic shear
  - Generally lack kink and other higher order terms



# Variety of Modes in Addition to KBM Contribute to Particle, Heat, Impurity Transport

#### Strong ongoing effort via DOE/FES milestones this year

 Electron heat transport driven by a range of instabilities driven by T<sub>e</sub> gradients and trapped particles

Eg, broad spectrum of electron drift waves in NSTX Enhanced Pedestal H-mode, ETG [Gerhardt14,Battaglia17,Guttenfelder19]

Microtearing may play a role near the top or inside the pedestal

Simulations of MAST, NSTX, JET [Dickinson13, Canik13, Hatch17, Battaglia17]

- Toroidal Ion Temperature Gradient (ITG) Modes Stable, but remnant slab-like ITG remains [Hatch, Kotschenreuther16]
  - Potentially important mechanism for particle pinch
- Higher excitation states of TEM/ITG, coupled modes [Pueschel17, Belli10]
- Different interactions with flow shear [Hatch17]



<u>Open issue</u>: Quantitative understanding of full range of instabilities that regulate particle, heat, momentum and impurity transport in the pedestal



23

### Including Cross Separatrix Geometry Important

1902-10299

- Important neoclassical effects associated with ion orbit loss near the separatrix
  - Key role in understanding E, and possibly L-H transition
  - Interaction with neutrals and sheath
- Electrostatic GK simulations and Electromagnetic GF simulations find strong turbulence, filaments ejected across separatrix [S. Ku/CS Chang 17, XQ Xu 17]





# Outline: Pedestal Physics Key to Predicting and Optimizing the Tokamak

- The Pedestal: What it is and why it matters
  - Simultaneous improvement of confinement and stability
  - Predictive capability enables fusion power optimization ( $P_{fus} \sim p_{ped}^2$ )
- Rich physics and computational challenges
  - Overlap of scales, challenge to methods (L~ $\lambda$ ~ $\rho$ )

#### Physics approaches and experimental tests

- Gyrokinetics and neoclassical theory
- MHD and peeling-ballooning modes
- Predicting and optimizing the pedestal: The EPED model
  - Development and testing
  - Coupled core-pedestal prediction fusion optimization
- Super H-Mode and high fusion performance
- ITER predictions and future prospects



## Large Pressure and Current Gradients in Pedestal Drive MHD Instabilities

#### • Potential Energy with stabilizing and destabilizing terms

- Negative energy implies MHD instability
- $\boldsymbol{\xi}$  = displacement of plasma fluid,  $\boldsymbol{B}_1$  = magnetic field perturbation

Compression of the magnetic field, (Fast, magneto-acoustic waves)

magnetic field line bending (Alfven waves)

compression (Slow, magneto-acoustic waves)

$$\delta W = \frac{1}{2} \int dV \left( \left| B_{1,\perp} \right|^2 + B_0^2 \left| \nabla \cdot \xi_{\perp} + 2\xi_{\perp} \cdot \kappa \right|^2 + \lambda p_0 \left| \nabla \cdot \xi \right|^2 \right)$$

$$-\int dV \left(2(\xi_{\perp} \cdot \nabla p_0)(\kappa \cdot \xi_{\perp}) + J_{0,\parallel}(\xi_{\perp} \times B_0 / B_0) \cdot B_{1,\perp}\right)$$

pressure gradient destabilizing (κ=field curvature) **ballooning** drive

parallel current destabilizing kink/peeling drive



# Peeling-Ballooning Modes Drive ELMs and Constrain the Pedestal Height



Pedestal is constrained, and ("Type I") ELMs triggered by intermediate wavelength (n~3-30) MHD instabilities called "peeling-ballooning" modes

•Driven by sharp pressure gradient and bootstrap current in the edge barrier (pedestal)

•Complex dependencies on  $v_*$ , shape etc., extensively tested against experiment

The P-B constraint is fundamentally non-local (effectively global on the scale of the barrier)

### Efficient MHD codes (eg ELITE, MISHKA, KINX) allow accurate computation of the intermediate n peeling-ballooning stability boundary enabling systematic comparison to observations

H.R. Wilson, P.B. Snyder et al PoP **9** 1277 (2002). P.B. Snyder, H.R. Wilson et al PoP **9** 2037 (2002), GTA Huysmans PPCF 47 (2005) B165, S. Saarelma et al PPCF 49 (2007) 31, M.G. Dunne et al PPCF 59 (2017) 025010, A. Merle et al PPCF 59 (2017) 104001 etc

27

### Peeling-Ballooning Model Successfully Applied Across a Range of Tokamaks

- International Tokamak Physics Activity (ITPA) coordinated multi-tokamak analysis
- Validated on all major international tokamaks
- ELM crash within 20% of calculated pedestal stability limit

Open issue: Recent JET metal wall cases sometimes exhibit ELMs below limit. Role of full rotation and extended MHD under investigation



DIII-D

### Observed ELM Spatial Structure Similar to Calculated Peeling-Ballooning Modes



[S. Pamela, Plasma Phys. Control. Fusion 55 095001 (2013)]

 Complicated structure but mode number similar to that calculated from linear stability



1902-10299

#### Nonlinear Simulations study ELM Dynamics and ELM Suppression/Control

- Nonlinear evolution of peeling-ballooning modes complex and can lead to bursting or saturated states
  - Saturated states (eg Quiescent H-Mode) are a promising technique for avoiding ELMs
  - Quantifying ELM losses is multi-scale, challenging. Being approached with extended MHD and gyrofluid techniques

#### • 3D perturbations can suppress and mitigate ELMs

Both linear response and nonlinear simulations exploring physics)

Open issues: Low resistivity leads to fine scale current sheets at computational resolution. Simulating a full ELM cycle with multi-scale physics is a grand computational challenge. Working models exist for RMP ELM suppression and QH mode but detailed quantitative understanding needed

#### JOREK: JET ELM Divertor



[G. Huijsmans, Nucl. Fusion (2013)]

#### **BOUT++ simulations**



# Outline: Pedestal Physics Key to Predicting and Optimizing the Tokamak

- The Pedestal: What it is and why it matters
  - Simultaneous improvement of confinement and stability
  - Predictive capability enables fusion power optimization ( $P_{fus} \sim p_{ped}^2$ )
- Rich physics and computational challenges
  - Overlap of scales, challenge to methods (L~ $\lambda$ ~ $\rho$ )
- Physics approaches and experimental tests
  - Gyrokinetics and neoclassical theory
  - MHD and peeling-ballooning modes
- Predicting and optimizing the pedestal: The EPED model
  - Development and testing
  - Coupled core-pedestal prediction fusion optimization
- Super H-Mode and high fusion performance
- ITER predictions and future prospects

31

### EPED Goal: Cut Through Complexity of Pedestal, Generate Predictive Model to Test and Improve

Paradigm: Transport barrier formation starts near separatrix and propagates inward primarily due to diamagnetic E<sub>r</sub>

Schematically divide instabilities that impact transport & stability in the pedestal into 2 categories: A. "Global" modes: extend across edge barrier including significant impact at top B. "Nearly-local" modes within the edge barrier

<u>Conjecture:</u> while neoclassical and electron microinstabilities drive transport, KBM commonly provides the final constraint on the pressure gradient.

- Key elements: neoclassical bootstrap current, nearly-local KBM, global peeling-ballooning
- Here take pedestal density as input
  - predicting it is a key goal for future work



#### **Mechanics of the EPED Predictive Model**

- Input: B<sub>t</sub>, I<sub>p</sub>, R, a,  $\kappa$ ,  $\delta$ , n<sub>ped</sub>, m<sub>i</sub>, [ $\beta$ <sub>global</sub>, Z<sub>eff</sub>]
- Output: Pedestal height and width (no free or fit parameters)
- A. P-B stability calculated via a series of model equilibria with increasing pedestal height
  - ELITE, n=5-30; non-local diamag model from BOUT++ calculations



P.B. Snyder et al Phys Plas 16 056118 (2009), NF 51 103016 (2011)



#### **Mechanics of the EPED Predictive Model**

- Input: B<sub>t</sub>, I<sub>p</sub>, R, a,  $\kappa$ ,  $\delta$ , n<sub>ped</sub>, m<sub>i</sub>, [ $\beta$ <sub>global</sub>, Z<sub>eff</sub>]
- Output: Pedestal height and width (no free or fit parameters)
- A. P-B stability calculated via a series of model equilibria with increasing pedestal height
  - ELITE, n=5-30; non-local diamag model from BOUT++ calculations

**B.** KBM Onset: 
$$\Delta_{\psi_N} = \beta_{p,ped}^{1/2} G(v_*, \varepsilon...)$$

 Directly calculate with ballooning critical pedestal technique



P.B. Snyder et al Phys Plas **16** 056118 (2009), NF **51** 103016 (2011)

• Different width dependence of P-B stability (roughly  $p_{ped} \sim \Delta_{\psi}^{3/4}$ ) and KBM onset ( $p_{ped} \sim \Delta_{\psi}^2$ ) ensure solution, which is the EPED prediction (black circle)



#### **Mechanics of the EPED Predictive Model**

- Input: B<sub>t</sub>, I<sub>p</sub>, R, a,  $\kappa$ ,  $\delta$ , n<sub>ped</sub>, m<sub>i</sub>, [ $\beta$ <sub>global</sub>, Z<sub>eff</sub>]
- Output: Pedestal height and width (no free or fit parameters)
- A. P-B stability calculated via a series of model equilibria with increasing pedestal height
  - ELITE, n=5-30; non-local diamag model from BOUT++ calculations
- **B.** KBM Onset:  $\Delta_{\psi_N} = \beta_{p,ped}^{1/2} G(v_*, \varepsilon...)$ 
  - Directly calculate with ballooning critical pedestal technique



P.B. Snyder et al Phys Plas 16 056118 (2009), NF 51 103016 (2011)

- Different width dependence of P-B stability (roughly  $p_{ped} \sim \Delta_{\psi}^{3/4}$ ) and KBM onset ( $p_{ped} \sim \Delta_{\psi}^{2}$ ) ensure solution, which is the EPED prediction (black circle)
  - can then be systematically compared to existing data or future experiments
- P-B stability and KBM constraints are tightly coupled: If either physics model (A or B) is incorrect, predictions for both height and width will be systematically incorrect

Effect of KBM constraint is counter-intuitive: Making KBM stability worse increases pedestal height and width

(eg "wide pedestal quiescent H-Mode,")

35

#### Numerous Experimental Tests of EPED Conducted



### Validation efforts coordinated with ITPA pedestal group, US JRT

- >800 Cases on 6 tokamaks
- Broad range of density (~1-24 10<sup>19</sup>m<sup>-3</sup>), collisionality (~0.01-4), f<sub>GW,ped</sub> (~0.1-1.0), shape (δ~0.05-0.65), q~2.8-15, pressure (1.7 - 35 kPa), β<sub>N</sub>~0.6-4 B<sub>t</sub>=0.7-8T
- Includes experiments where predictions were made before expt
- Typical σ~20-25%
- Recent work on TCV (Merle, Sauter, Medvedev PPCF17, Sheikh et al PPCF19 etc)



#### Outline: Pedestal Physics Key to Predicting and Optimizing the Tokamak

- The Pedestal: What it is and why it matters
  - Simultaneous improvement of confinement and stability
  - Predictive capability enables fusion power optimization ( $P_{fus} \sim p_{ped}^2$ )
- Rich physics and computational challenges
  - Overlap of scales, challenge to methods (L~ $\lambda$ ~ $\rho$ )
- Physics approaches and experimental tests
  - Gyrokinetics and neoclassical theory
  - MHD and peeling-ballooning modes

#### • Predicting and optimizing the pedestal: The EPED model

- Development and testing
- Coupled core-pedestal prediction fusion optimization
  - Skipping in interest of time: Combining EPED with core transport models such as TGLF+NEO enable prediction and optimization of global confinement
- Super H-Mode and high fusion performance
- ITER predictions and future prospects

37

## Outline: Pedestal Physics Key to Predicting and Optimizing the Tokamak

- The Pedestal: What it is and why it matters
  - Simultaneous improvement of confinement and stability
  - Predictive capability enables fusion power optimization ( $P_{fus} \sim p_{ped}^2$ )
- Rich physics and computational challenges
  - Overlap of scales, challenge to methods (L~ $\lambda$ ~ $\rho$ )
- Physics approaches and experimental tests
  - Gyrokinetics and neoclassical theory
  - MHD and peeling-ballooning modes
- Predicting and optimizing the pedestal: The EPED model
  - Development and testing
  - Coupled core-pedestal prediction fusion optimization
- Super H-Mode and high fusion performance
  - Can we use what we've learned to do more than just understand existing regimes?
- ITER predictions and future prospects



### Thinking Outside the Box: Super H Mode

- EPED model normally predicts a single pedestal solution
- At strong shaping, fixed input parameters (including density), PB mode can go from stable to unstable (pressure driven) and back to stable again with increasing pressure and current: multiple roots for two "equations", PB and KBM
- Expect only lowest solution to be accessible for these parameters. However, can move in third dimension (eg density) to access higher roots (Super H)





### At High Density and Strong Shaping, Solution Splits into H-Mode and Super H



#### • Constant density trajectories lead to usual H-Mode solution

- Optimal density leads to high pedestal near Super H (blue)
- Solution above H-mode (red) called Super H-Mode
  - Much higher pedestal than equivalent H-Mode solution



### At High Density and Strong Shaping, Solution Splits into H-Mode and Super H



 Super H-Mode Regime can be reached by dynamic optimization of the density trajectory

- Start at low density, and increase density over time (red arrow).
- Very high Super H-Mode pedestal should enable both high confinement and higher beta limit (broader profiles), leading to high fusion performance



41

### Super-H Mode Regime Accessed on DIII-D

EPED Predictions Compared to DIII-D Observations t=3.7s 25 t=3.5s Pedestal Height (kPa) t=2.9s 20 t=2.2s = W. Solomon PRL 113 135001 (2014) 15 t=1.7s t=4.3s P.B. Snyder NF 55 083026 (2015) 10 Super H-Mode H-Mode Near Super H 0 2 З 6 7 8 9 10 4 Pedestal Density \* Sqrt(Z<sub>eff</sub>/2.7) (10<sup>19</sup> m<sup>-3</sup>)

- Very high  $p_{ped}$  reached in density ramp with strong shaping ( $\delta \sim 0.53$ )
- Good agreement with EPED, which predicts this is the Super-H regime for n<sub>eped</sub>>~5.5
- Clear indication of bifurcation in p<sub>ped</sub>(n<sub>eped</sub>)
- Super H regime accessed sustainably with quiescent edge

# High peak performance in Super H-Mode experiments



#### Very High Super H Mode Pressure Predicted for C-Mod



- Alcator C-Mod is a compact, high field device (here  $B_t \sim 5.3T$ ), capable of high  $\delta$ 
  - After discovery of Super H-Mode on DIII-D, predictions were made for C-Mod (right)
    - Test SH theory at high  $B_t \& B_p$ , zero injected torque (RF), high Z metal wall
  - Following the right parametric trajectory should enable very high pressure
    - Need to reach densities much lower than typical for C-Mod H-mode to access Super H
    - Challenging to do on a high-Z metal wall device like C-Mod

Alcatol C-Mod



#### Access to Super H Mode on C-Mod Achieved via L-I-H Transition



- Transitioning first to I-mode, then to H-mode leads to a low n<sub>e</sub>, low impurity H-mode (left)
- As pedestal approaches predicted kink/peeling limit, low n mode observed (center)
- Discharges at 1MA, 5.4T reach SH regime, p<sub>ped</sub>~70 kPa (right)





### Super H-Mode Experiments on C-Mod Yield ITER-like pped



 Super H-Mode expt at 1.4MA achieved record 81 kPa pedestal pressure on last day of Alcator C-Mod operations, ITER-like pressure at ITER-like field [Hughes NF 2018]

- EPED model successfully tested over 2 orders of magnitude in pressure on 6 tokamaks
  - No indication of significant variation of model accuracy with  $\rho^*$  or  $p_{\text{ped}}$

Alcator C-Mod





#### Broad Profiles and High Pressure Obtained in Both C-Mod and DIII-D



- High pedestal pressure enables good confinement, high global MHD limits
  - C-Mod:  $B_{t}$ =5.3-5.8T,  $I_{p}$ =0.8-1.4MA, a=0.19m, R=0.67m,  $\delta$ ~0.5
    - ~ 100-170 kPa, p<sub>ped</sub> ~ 50 80 kPa

Alcator C-Mod

- DIII-D:  $B_t=2.1-2.2T$ ,  $I_p=1.6-2.0MA$ , a=0.6m, R=1.67m,  $\delta$ ~0.5-0.7
  - \_ ~ 70 110 kPa, p<sub>ped</sub> ~ 20 32 kPa, T<sub>i,0</sub> ~ 14 18 keV



## Very High Pedestal Pressure, Stored Energy, and Confinement Time in Recent co-I<sub>p</sub> Super H-Mode Experiments on DIII-D



- Deep access into Super-H regime, good agreement with EPED predictions
  - B<sub>t</sub>=2.17T, I<sub>p</sub>=1.6-2.0MA, a=0.6m,  $\delta$ ~0.5-0.7
  - p<sub>ped</sub>~30kPa, W~2 3.2 MJ (highest in present DIII-D config.) at modest P<sub>nbi</sub> ~ 8-12 MW
  - Peak t~0.4-0.7s,  $H_{98}$ ~2.2-2.9, t<sub>E</sub> ~ 30-67 kPa s, nTt ~ 4 8 10<sup>20</sup> keV m<sup>-3</sup> s



#### High Pedestal Pressure and T<sub>i,ped</sub> Enable High Peak Fusion Performance on DIII-D



• DD neutron rates up to 1.85 10<sup>16</sup>/s

- ~2/3 thermal, P<sub>fus,DD</sub> ~ 22 kW, P<sub>fus,DT,eq</sub> ~ 4.8 MW (at P<sub>nbi</sub> ~ 9 MW)



49

#### High Pedestal Pressure and T<sub>i</sub> Enable High Peak Fusion Performance on DIII-D, Record Fusion Gain



- Equivalent  $Q_{DT,eq} = P_{fus,DTeq}/P_{nbi} \sim 0.54$ .  $Q_{DT,eq}^* = P_{fus,DTeq}/(P_{nbi}-dW/dt) \sim 1$ 
  - Previous DIII-D record Q = 0.32, Lazarus96 in negative central shear discharges with 2.2MA,  $22m^3$
  - Achieved at modest B = 2.17T,  $I_p$ =2MA, V=20 m<sup>3</sup>. DT<sub>eq</sub> Fusion power density ~0.2 MW/m<sup>3</sup>

Appears to be highest  $Q_{DT,eq}$  and T on any medium size (R<2m) tokamak, and highest  $Q_{DT,eq}/IaB$  or  $Q_{DT,eq}/R^2B^2$  on any MFE device





## Super H-Mode Sustained Using 3D Magnetic Perturbations to Control Density and Impurity Accumulation

#### High performance condition sustained by applying 3D β<sub>N</sub> 2 magnetic perturbation Controls density and impurity accumulation 0

- Feedback control of pedestal or average density demonstrated
- Sustained W~1.9MJ,  $Q_{\text{DT,eq}}$ ~ 0.15,  $_{\text{T}}$ ~0.2s, H98~1.6,  $~\beta_{\text{N}}$ ~ 2.9
- ~2s sustainment (hardware limited)





## Predictions for ITER, Implications for Compact, High Performance Fusion



#### Dependence on $\rho^*$ Important for Predictions of ITER



- Key dimensionless parameters for ITER or DEMO reactor matched on existing machines (v\*,β,q, ε) except ρ\*
- Argument based on global ExB stabilization of turbulence leads to ρ\* dependence (γ~c<sub>s</sub>/L, ω<sub>E</sub>~p/L<sup>2</sup>, ω<sub>E</sub>>γ -> L < cρ )</li>
- EPED predicts ~no ρ\* dependence: front propagation model for barrier formation and broadening
- Observations find little/no p\* dependence, including JET metal wall [Beurskens, Osborne PPCF09, Maggi17]

#### <u>Open issue:</u> Important to continue testing and developing understanding at very small p\*



53

## ITER Pedestal Predictions Made for more than 15,000 cases, used to train neural net (ITPA)



- Fixed: R=6.2m, a=2m
- Varied:  $\kappa$ =1.7-1.9,  $\delta$ =0.45-0.49,  $\beta_N$ =1-3,  $Z_{eff}$ =1-3,  $m_i$ =1-3
- Three categories: full (5.3T), half (2.6T), and 1/3 (1.8T) field
  - Full field: I<sub>p</sub>=5-17MA (most 7-15MA), n<sub>eped</sub>=3-15 10<sup>19</sup> (most 6-10.6)
  - Half field: Ip=2-10MA (most 7.5MA), neped=3-10.6 10<sup>19</sup>
- 6 dimensional scan at each of 3 B<sub>t</sub> values: used to train neural net

GENERAL

## Super H/NSH Regime Access is Predicted for ITER: DIII-D has Achieved Needed $\beta_{N,ped}$ , $n_{e,sep}$ , $n_{e,ped}$ Consistently





Open issue: Physics of the Greenwald density limit which constrains degree of Super H access and predicted performance for ITER and DEMO concepts

- Core-pedestal simulations find ITER high performance (Q>10) at high n<sub>e</sub> [Meneghini16]
- DIII-D SH experiments reproduce many characteristics of the predicted ITER regime, including β<sub>N,ped</sub>~0.8, n<sub>e,sep</sub>~3-4, n<sub>e,ped</sub>~7-10. C-Mod produces p<sub>ped</sub>~80 kPa
  - Potential for substantial improvements in ITER performance, consistent with n<sub>e,sep</sub>

NERAL ATOMICS

#### Super H and Near Super H Operation Enables Very High Fusion Performance per I<sub>p</sub>aB<sub>t</sub>



#### Simple metric of fusion performance (Q or W/P) per l<sub>p</sub>aB<sub>t</sub>

- Colored points are observations ( > 50 kPa), red points are SH/NSH experiments
- High Q/IaB enables ITER success, and compact, cost attractive pilot plant

**GENERAL ATOMICS** 

#### Summary: Exciting New Discoveries in Pedestal Physics Leading to Improvements in Fusion Performance

- Multi-scale nature of pedestal leads to rich physics that challenges traditional analytic
   & computational approaches
  - Open issues: formalism, particle & momentum transport, impurities, neutrals, ρ\*, L-H, i-mode & WPQH...
- Despite challenges, significant progress made via gyrokinetic/neoclassical, MHD approaches, combined with advanced diagnostics
  - Extensive validation studies on flexible tokamaks with high resolution measurements
- Simple model (EPED) predicts pedestal height to ~20-25% accuracy in many regimes. Coupling to core models enables initial global confinement prediction
  - Revolutionary capability for tokamak fusion optimization (many open issues, connection to SOL)
  - Gyrokinetic/neoclassical studies working toward predictive capability for individual transport channels (n,T,v)

#### • Super H regime enables high pedestal and high fusion performance

- Predictions guided experiments, leading to discovery of new regime
- Record ITER-like pedestal pressure on Alcator C-Mod, high fusion performance on DIII-D (Q<sub>DT,eq</sub>~ 0.5)
- Potential for high performance in ITER, and compact, high performance fusion reactors
  - ITER predictions made for >15,000 cases, used to train neural net for efficient testing and coupled corepedestal simulations. Developing accurate methods for incorporating SH solutions as well.

![](_page_56_Picture_13.jpeg)

#### Acknowledgments

- This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Fusion Energy Sciences, under Awards DE-FC02-04ER54698, DE-FC02-99ER54512, DE-FG02-95ER54309, DE-FC02-06ER54873, DE-SC0014264, and DE-SC0017992, using the DIII-D National Fusion Facility and Alcator C-Mod, DOE Office of Science user facilities. Extensive discussions with the ITPA pedestal group, as well as the numerous contributions of the Alcator C-Mod team, DIII-D team, and OMFIT group are gratefully acknowledged.
- DISCLAIMER: This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

![](_page_57_Picture_3.jpeg)

# **Extra Slides**

![](_page_58_Picture_1.jpeg)

### Neoclassical Theory Describes Collisional Transport Associated with Poloidal 'Banana' Orbits

![](_page_59_Figure_1.jpeg)

Collisions along orbits drive ion heat & momentum and impurity transport

RAL ATOMICS

### Neoclassical Physics also leads to Large "Bootstrap" Current in Pedestal

Gyro-orbits drift due to non-uniform field  $\Rightarrow$  banana orbits

![](_page_60_Figure_2.jpeg)

But <u>more & faster</u> particles (strong density and temperature gradients) on orbits nearer the core (green cf blue) lead to a net "banana current"

- this is transferred to a helical bootstrap current via collisions

1902-10299 – Bootstrap current typically dominant in pedestal, major role in instabilities

# Sustainment and Core-Edge Compatibility of Super H-Mode Regime

![](_page_61_Picture_1.jpeg)

### Connecting a High Performance Super H Pedestal & Core to a High Density, Radiative Divertor & SOL

- Super H (J-limited) solution predicted not to show degradation of pedestal pressure w/ n<sub>e,sep</sub>
  - P-limited solution degrades with increasing  $n_{e,\text{ped}}$  and  $n_{e,\text{sep}}$  (eg high gas puff in JET ILW)
- Scan D<sub>2</sub> gas rate, and introduce radiative impurities (N<sub>2</sub>) into the Div/SOL to test predictions on DIII-D
  - Use 3D magnetic perturbations (i-coil) to control particle and impurity accumulation in core
  - Use i-coil feedback to maintain ~constant density in pedestal
     & core as separatrix, divertor and SOL density are increased
    - Test EPED predictions of sensitivity of pedestal to separatrix conditions

![](_page_62_Figure_7.jpeg)

![](_page_62_Figure_8.jpeg)

![](_page_62_Picture_9.jpeg)

![](_page_62_Picture_10.jpeg)

#### D<sub>2</sub> gas Scan Increases Separatrix and Divertor Density while Pedestal Pressure and Confinement Remain High

- D<sub>2</sub> gas scan in Super H mode experiment at I<sub>p</sub>=2MA, B<sub>t</sub>=2.1T. Gas rate varied ~30x
  - Pedestal pressure and  $\tau_E$  remain ~fixed, high
  - i-coil feedback control of  $n_{e,ped} \sim 7\text{-}8~10^{19}~m^{\text{-}3}$  successful up to~110 torrL/s of  $D_2$  gas
  - Separatrix density rises from  $\sim 2.5 4 \ 10^{19} \ m^{-3}$
  - Strike point density rises from  $\sim 2.5 7 \ 10^{19} \text{ m}^{-3}$

### Both pedestal and separatrix density reach ITER values while maintaining high confinement and $p_{ped}$

Super H-mode compatible with both high fusion performance and high separatrix density for divertor solutions.

![](_page_63_Picture_8.jpeg)

![](_page_63_Figure_9.jpeg)

#### N<sub>2</sub> Injection Effective for Cooling Divertor while Maintaining High Performance Core & Pedestal

- Significant cooling with ~5MW of divertor radiated power using feedback on N<sub>2</sub>
  - Peak T<sub>e</sub> near strike point drops more than 3x
  - Pedestal pressure and confinement remain ~constant
  - Future experiments needed to explore full detachment and impact of closed divertor

![](_page_64_Figure_5.jpeg)

65

![](_page_64_Picture_6.jpeg)

## Integrated Modeling Enables Prediction and Optimization of Coupled Core-Pedestal System

- Peeling-ballooning stability is enhanced by the global Shafranov shift, which is proportional to global pressure [Snyder07,Chapman15,Saarelma17]
- Core turbulent transport is gradient scale length driven, and hence core profiles depend strongly on the BC provided by the pedestal
- Potential for a virtuous cycle to strongly enhance performance, but must do selfconsistent, coupled pedestal-core modeling

![](_page_65_Figure_4.jpeg)

![](_page_65_Figure_5.jpeg)

![](_page_65_Picture_6.jpeg)

### Example: EPED/TGLF/NEO and Core-Pedestal Integrated Modeling: DIII-D ITER-similar Discharge 153523

- Divide plasma into 4 regions
- Coupled workflow with OMFIT/IPS

![](_page_66_Figure_3.jpeg)

![](_page_66_Figure_4.jpeg)

### Example: EPED/TGLF/NEO and Core-Pedestal Integrated Modeling: DIII-D ITER-similar Discharge 153523

- No measurements of T<sub>e</sub>, T<sub>i</sub> or pressure input
- Density only input at pedestal
  - Inputs: shape, sources, rot.,  $B_t$ ,  $I_p$ ,  $n_{e,ped}$
  - Predicting  $T_e,\,T_i,\,n_{e,core},\beta_N$
- Step 1: Run EPED
  - Don't yet know  $\beta_{N}$  so use (poor) initial guess
- Step 2: Run TGYRO using BC from EPED to predict profiles and  $\beta_{\text{N}}$
- Step 3: Run EPED using updated value for  $\beta_N$
- •••
- Iterate to convergence
  - Have predicted profiles for  $T_e,\,T_i,\,n_e$  and pressure/ $\beta_N$
  - Result independent of initial guess

![](_page_67_Figure_13.jpeg)

### Example: EPED/TGLF/NEO and Core-Pedestal Integrated Modeling: DIII-D ITER-similar Discharge 153523

- Accurately predicts full T<sub>i</sub> and T<sub>e</sub> profile, core density profile and global beta in this case
  - Core-pedestal coupling essential to achieve this
  - Statistical accuracy in large studies

#### Revolutionary capability

- Predict confinement and stored energy without empirical scalings
- Employing to predict and optimize the performance of ITER and future devices

<u>Open issues</u>: Predicting L mode and L-H transition. Coupling to open field line region and divertor, material surfaces. Predicting particle and impurity transport through pedestal.

![](_page_68_Figure_8.jpeg)

#### Super H-Mode Experiments on Alcator C-Mod and DIII-D Achieve High Fusion Performance, Record Pedestal Pressure

- Super H-mode (SH) predicted in strongly shaped plasmas: high pped, increases with ne [Snyder NF15]
- Record pedestal pressures (~80 kPa) achieved in C-Mod SH experiments [Hughes NF18]
  - Successful tests of EPED model up to ~90% of predicted ITER p<sub>ped</sub>

Alcator

C-Mod

- **Record DIII-D fusion gain** ( $Q_{DI,eq} \sim 0.54$ ).  $Q_{DI,eq}/IaB$  and  $Q_{DI,eq}/(RB)^2$  highest reported on any tokamak •
- High performance sustained w/ 3D magnetic perturbations to control  $n_e$  and impurity accumulation •
- Predicted to enable high performance on ITER, and be compatible with high separatrix density for divertor solutions

![](_page_69_Figure_7.jpeg)

![](_page_69_Picture_8.jpeg)

70